

A presenilin-1-dependent γ -secretase-like protease mediates release of Notch intracellular domain

Bart De Strooper*, Wim Annaert*, Philippe Cupers*, Paul Saftig†, Katleen Craessaerts*, Jeffrey S. Mumm‡, Eric H. Schroeter‡, Vincent Schrijvers*, Michael S. Wolfe§, William J. Ray||, Alison Goate|| & Raphael Kopan‡

* Neuronal Cell Biology and Gene Transfer Laboratory, Flanders Institute for Biotechnology (VIB4), Center for Human Genetics, KU Leuven, B-3000 Leuven, Belgium

† Zentrum Biochemie und Molekulare Zellbiologie, Abteilung Biochemie II, Universität Göttingen, B-37073, Germany

‡ Division of Dermatology and the Department of Molecular Biology and Pharmacology, Washington University, St Louis, Missouri 63110, USA

§ Department of Pharmaceutical Sciences, University of Tennessee, Memphis, Tennessee 38138, USA

|| Departments of Psychiatry and Genetics, Washington University School of Medicine, St Louis, Missouri 63110, USA

Signalling through the receptor protein Notch, which is involved in crucial cell-fate decisions during development, requires ligand-induced cleavage of Notch. This cleavage occurs within the predicted transmembrane domain, releasing the Notch intracellular domain (NICD), and is reminiscent of γ -secretase-mediated cleavage of β -amyloid precursor protein (APP), a critical event in the pathogenesis of Alzheimer's disease. A deficiency in presenilin-1 (PS1) inhibits processing of APP by γ -secretase in mammalian cells, and genetic interactions between Notch and PS1 homologues in *Caenorhabditis elegans* indicate that the presenilins may modulate the Notch signalling pathway¹⁻⁴. Here we report that, in mammalian cells, PS1 deficiency also reduces the proteolytic release of NICD from a truncated Notch construct, thus identifying the specific biochemical step of the Notch signalling pathway that is affected by PS1. Moreover, several γ -secretase inhibitors block this same step in Notch processing, indicating that related protease activities are responsible for cleavage within the predicted transmembrane domains of Notch and APP. Thus the targeting of γ -secretase for the treatment of Alzheimer's disease may risk toxicity caused by reduced Notch signalling.

Notch-1 is synthesized as a type I integral-membrane protein of relative molecular mass 300,000 (M_r 300K) (Fig. 1) that is cleaved by a furin-like protease in the Golgi during trafficking of Notch to the cell surface. The two proteolytic fragments remain associated to form the functional receptor^{5,6}. Following ligand binding, Notch-1 undergoes further cleavage close to or within its transmembrane domain, releasing the NICD. The NICD translocates to the nucleus and modifies transcription of target genes through its association with CSL proteins (where CSL stands for CBF1, Su(H), Lag-1)⁷⁻⁹.

APP is also a type I integral-membrane protein. APP is the precursor of the amyloid- β peptide, which is deposited in the brains of Alzheimer's disease patients. Amyloid- β peptide is derived from APP as a result of cleavage of APP by β - and γ -secretase, whose molecular nature remains unclear. β -Secretase cleaves the extracellular domain of APP, producing a soluble ectodomain and a membrane-associated carboxy-terminal fragment (Fig. 1). γ -Secretase catalyses an intramembranous cleavage of this membrane-associated fragment, resulting in generation of amyloid- β peptide and the production of a C-terminal fragment of APP. Cleavage of APP by α -secretase precludes production of amyloid- β peptide (Fig. 1). The




Figure 1 Proteolytic cleavage of Notch, APP and SREBP (not to scale). The proteolytic fragments (NICD and α - and β -stubs) and constructs (mN Δ E) used to analyse Notch-1 and APP metabolism are indicated, as are the proteases involved in processing (furin, γ' -, α -, β - and γ -secretases, and site-1 and site-2 proteases). For a more extensive discussion on the analogies between APP, Notch-1 and SREBP processing, see ref. 30.

functional role(s) of these APP fragments is unknown. Genetic and biochemical studies have implicated the proteolytic processing events that lead to amyloid- β -peptide generation in the pathogenesis of Alzheimer's disease. Mutations in APP and the presenilins (PS1 and PS2) cause familial early-onset Alzheimer's disease, and presenilin mutations lead to increased processing of APP by γ -secretase *in vitro* and *in vivo*¹⁰. PS1-deficient mice show decreased γ -secretase processing of APP⁴ and developmental abnormalities consistent with altered Notch signalling^{2,3}. Genetic interactions between the *notch* homologues *glp-1* and *lin-12* and the *presenilin* homologues *sel-12* and *hop-1* in *C. elegans* provide indirect evidence for the involvement of the presenilins in the Notch signalling pathway¹¹. Here we identify the specific step in Notch signalling that is controlled by PS1.

We first investigated whether the absence of PS1 interferes with the normal expression of Notch, as has been suggested². We used brain extracts from PS1-deficient (PS1^{-/-}) mouse embryos at embryonic day (E) 14. As expected, no expression of the amino-terminal or C-terminal fragments of PS1 was found (Fig. 2a and ref. 4). Moreover, in agreement with our previous results obtained from study of neuronal cultures⁴, we observed an accumulation of endogenous mouse APP C-terminal fragments in the same brain extracts, indicating deficient γ -secretase processing of APP when PS1 is absent (Fig. 2a, lower arrow). These fragments are predominantly 'stubs' produced by α -secretase, because endogenous mouse APP is not a good substrate for β -secretase (owing to the presence of amino-acid substitutions between the mouse and the human sequences at the β -secretase site)^{4,12}. In the same homogenates, we detected normal steady-state levels of furin-processed Notch-1 C-terminal fragments, indicating that PS1 is not required for Notch biosynthesis or cleavage by furin within the Golgi (Fig. 2b). PS1-independent Notch maturation was also observed *in vitro*, in mouse fibroblasts derived from PS1^{-/-} embryos (Fig. 2c). We conclude that PS1^{-/-} mouse cells produce normal amounts of mature (furin-cleaved) Notch 1 protein.

The similarities between the proteolytic processing of Notch and APP are striking, especially the final proteolytic events that release the cytoplasmic domains, raising the possibility that PS1 might mediate the intramembranous proteolytic cleavage of Notch-1 to




Figure 2 Processing of APP and Notch-1 in PS1^{-/-} mouse brain and MEFs. Proteins were detected in western blots. **a**, Brain-membrane extracts (4–20% SDS-PAGE, 50 μg per lane) were reacted with APP675–695, an antiserum specific for the APP C terminus, or with an antiserum specific to the PS1 N terminus (PS1NTF). Arrows identify PS1NTF and a mouse APP fragment generated by α-secretase-mediated processing (mouse APP α-stubs). **b**, Notch-1 was

immunoprecipitated from 100 μg brain extract using IC-antiserum, an anti-Notch-1 antiserum⁵ and stained with the monoclonal antibody Tan20 (ref. 6) (6% SDS-PAGE). TMIC identifies the furin-generated mature Notch-1 fragment. **c**, MEF cell extracts were reacted with mN1A, a monoclonal antibody against Notch-1 (4–20% SDS-PAGE, 50 μg per lane).

release NICD^{7–9}. As endogenous NICD levels in cells and tissues are biochemically below the level of detection⁷, we took advantage of two well-characterized Notch constructs, mNotchΔE and N^{ICV}, that have been used to demonstrate the relationship between processing of and signalling through Notch-1 (refs 7, 13). The mNotchΔE construct codes for the transmembrane and intracellular domains of murine Notch-1 and undergoes a single proteolytic cleavage to release NICD. This cleavage resembles cleavage of endogenous Notch-1 in all respects^{7,13}. The N^{ICV} construct encodes the intracellular domain of Notch-1 without the membrane anchor, producing a protein identical to NICD. N^{ICV} has been used as a control for the fate of NICD in PS1^{-/-} cells⁷. Both constructs have part of their C-terminal domain replaced with six consecutive Myc-epitope tags⁷. We cloned these constructs into recombinant Semliki forest virus (SFV) vector, to enable their expression in primary cultures of neurons⁴. The expression and subcellular localization of mNotchΔE were similar in wild-type and PS1^{-/-} cells, as was also the case for N^{ICV} (Fig. 3). However, NICD protein produced from N^{ICV} translocated to the nucleus in both PS1^{+/+} and PS1^{-/-} neurons (Fig. 3, lower panels), whereas nuclear staining in wild-type and PS1^{-/-} neurons expressing mNotchΔE was almost undetectable, indicating that the relative amounts of NICD compared with unprocessed mNotchΔE were below the detection limit of our immunofluorescence assay (Fig. 3, upper panels). These observations are in agreement with genetic evidence from *C. elegans* that *sel-12* mutations do not interfere with *lin-12* signalling when the LIN-12 intracellular domain is expressed alone¹¹.

To assay directly the role of PS1 in mNotchΔE cleavage, we performed pulse-chase experiments and analysed the generation of NICD. Within 30–60 minutes after labelling with ³⁵S-methionine, NICD was first detected in PS1^{+/+} cultures. NICD continued to accumulate throughout the 120-minute chase period (Fig. 4a, b; ratio of NICD to total mNotchΔE present at the start point of the chase). In contrast, NICD production was greatly reduced in PS1^{-/-} neurons (Fig. 4a, b). As the PS1^{-/-} embryos tend to yield fewer cells than their wild-type littermates⁴, we analysed twice the amount of PS1^{-/-} cell extracts. Only residual amounts of NICD were detected in PS1^{-/-} neurons (Fig. 4c), even when the mNotchΔE signal was stronger than in the PS1^{+/+} cell extracts. This difference could not be explained by a higher turnover rate of NICD in PS1^{-/-} neurons, as NICD produced from control N^{ICV} constructs showed the same stability in both PS1^{-/-} and PS1^{+/+} cells (Fig. 4d).

To test the generality of this observation in another, non-neuronal, cell type, we transiently transfected PS1^{-/-} fibroblasts with mNotchΔE plasmids. Although NICD was produced in PS1^{+/+}




Figure 3 Nuclear transport of the intracellular domain (NICD) of Notch-1. Neurons were transfected with SFV driving expression of mNotchΔE or N^{ICV} (which is identical to the NICD), fixed and immunostained. Red fluorescence indicates Myc-tagged mNotchΔE (upper panels) or NICD fragment (lower panels). Nuclear import of NICD is not disturbed in the PS1^{-/-} neurons. The green fluorescence in the lower panels indicates immunoreactivity towards calnexin, revealing the ER. The presence of uninfected neurons in the lower panels shows the specificity of the anti-Myc staining. +/+, PS1^{+/+} neurons; -/-, PS1^{-/-} neurons.

cells, it was hardly detectable in PS1^{-/-} cell extracts (Fig. 4e, extract). Co-immunoprecipitation of NICD with its nuclear partner, Flag-tagged CSL^{RBp3} (ref. 7), confirmed the existence of very low amounts of NICD in the PS1^{-/-} cells (Fig. 4e). As residual amounts of both NICD and amyloid-β peptide can be detected in PS^{-/-} cells, it is possible that the PS1 homologue PS2 (expressed in PS1^{-/-} cells; K.C. and B.D.S., unpublished observations), can partially compensate for the loss of PS1 function. We conclude that NICD production is decreased markedly in PS1^{-/-} cells.

The site of cleavage of Notch-1, releasing NICD, may occur at an analogous position to the site of cleavage of APP by γ-secretase¹⁴, N-terminal to the RKRRRQ stop-transfer signal⁷. This apparently




Figure 4 Generation of NICD in PS1^{+/+} and PS1^{-/-} neurons. **a**, Neurons transfected with SFV encoding mNotchΔE were labelled for 30 min and then chased in normal medium as indicated. Immunoprecipitated material was resolved in 4–20% SDS-PAGE. **b**, Phosphorimaging analysis of experiments performed as in **a**. The mNotchΔE signal at time zero was taken as 100% (lanes labelled 0' in **a**). NICD produced from mNotchΔE at each time point was normalized to this signal (produced fraction). Values are means ± s.e.m. (*n* = 3). **c**, Control experiment, performed as in **a** but using twice as much material from PS1^{-/-} neurons. NICD is hardly detectable at 120 min or 240 min of chase in PS1^{-/-} neurons (left arrow) even

when more mNotchΔE precursor is loaded. **d**, Turnover of NICD is identical in PS1^{+/+} and PS1^{-/-} neurons transfected with SFV encoding N^{lcv}. The neurons were labelled for 30 min followed by a chase as indicated. **e**, mNotchΔE was expressed in PS1^{+/+} and PS1^{-/-} MEFs. We then analysed MEF extracts (left) and immunoprecipitates produced using anti-Flag-CSL antibodies (right) by western blotting. NICD is present in PS1^{+/+} cell extracts and undetectable in PS1^{-/-} cell extracts. Enriching for NICD using anti-CSL co-immunoprecipitation reveals residual amounts of NICD in PS1^{-/-} cells (5 min exposure, compared to 20 s in the other lanes). Asterisk indicates a post-lysis degradation peptide.

intramembranous cleavage is unusual and occurs in only one other known protein, the sterol-regulatory-element-binding protein (SREBP)¹⁵ (Fig. 1). However, mutants deficient in SREBP cleavage show no abnormality in generation of amyloid-β peptide, indicating that the enzyme that mediates SREBP cleavage may not be closely related to γ-secretase¹⁶. No effects on SREBP processing were observed in PS1^{-/-} cells (data not shown). In contrast, our results from PS1^{-/-} cells show that PS1 is important for cleavage of both Notch-1 and APP. To determine whether the enzyme responsible for NICD release has similar pharmacological properties to γ-secretase, we compared the effects of protease inhibitors on processing of Notch-1 and APP, selecting inhibitors that have been shown previously to block γ-secretase^{17–20}. These inhibitors also interfere with mNotchΔE processing (Fig. 5 and refs 7, 13). MDL 28170, which is a peptide aldehyde calpain inhibitor, was the first inhibitor shown to interfere with γ-secretase¹⁸, and MG132, another peptide aldehyde, also inhibits γ-secretase. However, MG132 (and possibly MDL 28170) are relatively broad-spectrum protease inhibitors that inhibit the proteasome. We ruled out the possibility that inhibition of the proteasome is involved in the observed effects of MDL 28170 and MG132 on Notch-1 processing by using lactacystin, a specific proteasome inhibitor, which has no effect on NICD generation (Fig. 5 and E.H.S. and R.K., unpublished observations). Likewise, lactacystin does not affect γ-secretase-mediated cleavage of APP²¹.

To further explore the apparent similarity between γ-secretase and the Notch-1 protease, we used the γ-secretase inhibitor MW167, a difluoro ketone peptide analogue designed on the basis of the primary sequence of the site of cleavage by γ-secretase in

APP¹⁷. This compound inhibited, in a concentration-dependent manner, γ-secretase-mediated cleavage of human APP expressed in neurons using recombinant SFV (Fig. 6), leading to the accumulation of APP C-terminal stubs and the inhibition of amyloid-β-peptide secretion into the medium. As both α- and β-secretase-processed fragments appeared in the cells (α- and β-stubs, Fig. 6) and as secretion of the APP ectodomain into the medium was unaffected (data not shown), neither α- nor β-secretase enzymatic activity is inhibited by MW167 (see also ref. 17). Processing of mNotchΔE was inhibited by MW167 at concentrations similar to those that inhibit APP processing (Fig. 6). The estimated half-maximal inhibitory concentration (IC₅₀) for MW167 (10 μM) in mNotchΔE processing is nearly identical to its IC₅₀ for APP¹⁷. MW167 is also a potent inhibitor of cathepsin D²²; however, using a series of related difluoro ketone analogues, we found that the effect of MW167 on APP processing is not a consequence of cathepsin-D inhibition²². The three inhibitors (MG132, MDL28170 and the chemically different MW167) all inhibit mNotchΔE processing and γ-secretase-mediated cleavage of APP to similar extents. Although neither γ-secretase nor the enzyme that cleaves mNotchΔE has been purified yet, our results are consistent with the hypothesis that these two proteolytic activities are closely related.

We have shown that efficient processing of Notch-1 to produce NICD requires PS1 and is inhibited by γ-secretase inhibitors. We showed this by using the mNotchΔE construct, a reliable model for intracellular processing of Notch-1 (ref. 7). Furthermore, the expression of endogenous Notch-1, its maturation by a furin-like enzyme, and the translocation of NICD to the nucleus are not affected by PS1 deficiency. Notch-1 and APP both exhibit a genetic




Figure 5 Effect of γ -secretase inhibitors on mNotch Δ E processing. Neurons were transfected with SFV encoding mNotch Δ E as in Fig. 4a, c. Label was chased for 4 h in the presence of 1% DMSO (control), MDL 28170 (100 μ M), lactacystin (100 μ M) or MG132 (20 μ M). The stronger signals in the lactacystin lane probably reflect stabilization of mNotch Δ E when the proteasome is inhibited.




Figure 6 Effect of γ -secretase inhibitor MW167 on processing of mNotch Δ E (upper panel) and APP (lower panels). SFV-infected neurons were pulse-labelled for 30 min and label was chased for 4 h in the presence of DMSO vehicle or the indicated concentrations of MW167. mNotch Δ E and NICD were immunoprecipitated using anti-Myc antibodies. APP stubs (reflecting α -secretase- and β -secretase-mediated processing of human APP) and amyloid- β peptide (A β peptide; 4 h labelling) were immunoprecipitated as described^{4,12}.

interaction with presenilins^{1-3,11} and are cleaved by a γ -secretase-like activity that requires PS1 protein to perform an apparently intramembranous cleavage. These results provide a molecular explanation for the widely reported genetic interaction between Notch-1 and PS1 (refs 1-3, 11) in different species.

There is also a direct physical interaction between Notch and PS1 (ref. 23), as there is between APP and PS1/2 (refs 24, 25). The function of PS1 in Notch/APP processing could be that of a 'facilitator', analogous to the role of the endoplasmic-reticulum (ER)-resident, multimembrane-spanning protein SCAP in SREBP¹⁵ processing. Such a facilitator could directly expose APP and Notch-1 to the γ -secretase. Alternatively, PS1 could indirectly regulate the correct trafficking of APP, Notch-1 and γ -secretase to an as-yet-undefined subcellular (micro) domain where cleavage occurs²⁶. However, our results are also consistent with a more direct role for PS1 as a regulatory or catalytic component of the protease(s) that cleave(s) Notch-1 and APP. In this case, PS1 would have a role similar to that of the ER-resident multimembrane-spanning site-2 protease²⁷ in SREBP processing (Fig. 1).

Finally, our results have significant implications for the potential use of γ -secretase inhibitors as drugs for treatment of Alzheimer's disease. The resulting inhibition of Notch-1 processing in the adult would be predicted to lead to immunodeficiency and anaemia because of the important function of Notch-1 in haematopoiesis throughout life^{28,29}. Thus, it may be necessary to develop inhibitors specific for γ -secretase-mediated cleavage of APP or to target other steps in the generation and deposition of amyloid- β peptide. □

Methods

The generation and characterization of PS1^{-/-} mice, the generation of neuronal cell cultures, the use of recombinant SFV, metabolic labelling, cell extraction and double immunoprecipitation were done as described^{4,12}. For pulse-chase experiments, cells from one embryo were seeded in a 12-well or 4-well cell-

culture plate (Nunc) pretreated with polylysine and serum. After infection with the appropriate recombinant SFV^{4,12}, neurons were pulse-labelled for 30 min with 200 μ Ci ³⁵S-methionine per ml, and chased for the indicated time in normal medium. Cells were processed as before¹³. Every lane (Figs 4a, c, d, 5, 6) shows the material obtained from one well, and signals were compared between littermate embryos. MG132 (Calbiochem), MDL 28170 (provided by B. Cordell) and MW167, each dissolved in dimethylsulphoxide (DMSO), were added at the end of the pulse labelling to the indicated final concentrations. DMSO in the cell cultures was 1%.

Mouse brain was homogenized in 5 mM Tris-HCl, pH 7.4, 250 mM sucrose, 1 mM EGTA, 5 mM EDTA, 1 μ g ml⁻¹ pepstatin A and 100 U ml⁻¹ aprotinin, and centrifuged at 500g for 10 min. The resulting supernatant was centrifuged at 100,000g for 60 min. The pellet was resuspended in PBS and protein concentration was measured (Biorad protein assay).

We plated 5 \times 10⁵ mouse embryonic fibroblasts (MEFs) immortalized with SV40 large T antigen in 100-mM plates and transfected them by the Ca₂PO₄/BBS method¹³ with both Flag-CSL^{RBP3} and mNotch Δ E. After lysis in co-immunoprecipitation buffer⁷ containing 200 mM KCl, 10% of the sample was used for crude extract analysis. The remainder was co-immunoprecipitated⁷ using anti-Flag antibody. The washed pellet was resuspended in 50 μ l Laemmli lysis buffer and resolved by 6% SDS-PAGE, transferred to nitrocellulose membrane and visualized using 9E10 anti-Myc antibody^{7,13}. For immunocytochemistry, neurons cultured on glass coverslips were fixed in 4% paraformaldehyde and permeabilized in ice-cold methanol and acetone. Antibodies were diluted in blocking buffer and cover slips were mounted in Mowiol (Calbiochem). Cells were analysed using a MRC1024 confocal microscope (Biorad).

The complementary DNAs coding for mNotch Δ E and N^{ICV7,13}, antiserum APF675-695 (ref. 12) used to immunoprecipitate the APP C-terminal stubs, and antiserum APF597-612 (ref. 12) used to immunoprecipitate amyloid- β peptide have been described. We used mNotch Δ E in which Met 1727 was mutated to Val to prevent alternative translation initiation. NICD production from mNotch Δ E is unaffected by this mutation, as shown previously^{7,13}. B19/2 was raised against amino acids 30-44 of PS1. IC-antiserum against the cytoplasmic domain of murine Notch-1 (ref. 15) was provided by A. Israël and used to immunoprecipitate endogenous Notch-1 from brain extracts. Rat monoclonal antibody Tan20 (provided by S. Artavanis-Tsakonas) was used for immunodetection of precipitated Notch-1 (ref. 6). mN1A, a monoclonal antibody specific to the cytoplasmic domain of Notch-1 (provided by L. Milner), was used for detection of mouse Notch-1 from MEF SDS extracts. Calnexin antiserum and the anti-Myc monoclonal antibody 9E10 were provided by A. Helenius and J. Creemers.

Received 5 November 1998; accepted 5 February 1999.

- Levitán, D. & Greenwald, I. Facilitation of lin-12-mediated signalling by sel-12, a *Caenorhabditis elegans* S182 Alzheimer's disease gene. *Nature* **377**, 351-354 (1995).
- Wong, P. C. et al. Presenilin 1 is required for Notch1 and Dll1 expression in the paraxial mesoderm. *Nature* **387**, 288-292 (1997).
- Shen, J. et al. Skeletal and CNS defects in presenilin-1-deficient mice. *Cell* **89**, 629-639 (1997).
- De Strooper, B. et al. Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* **391**, 387-390 (1998).
- Logeat, F. et al. The Notch1 receptor is cleaved constitutively by a furin-like convertase. *Proc. Natl Acad. Sci. USA* **95**, 8108-8112 (1998).
- Blaumueller, C. M., Qi, H., Zagouras, P. & Artavanis-Tsakonas, S. Intracellular cleavage of Notch leads to a heterodimeric receptor on the plasma membrane. *Cell* **90**, 281-291 (1997).
- Schroeter, E. H., Kisslinger, J. A. & Kopan, R. Notch-1 signalling requires ligand-induced proteolytic release of intracellular domain. *Nature* **393**, 382-386 (1998).
- Struhl, G. & Adachi, A. Nuclear access and action of notch *in vivo*. *Cell* **93**, 649-660 (1998).
- Lecourtis, M. & Schweisguth, F. Indirect evidence for Delta-dependent intracellular processing of notch in *Drosophila* embryos. *Curr. Biol.* **8**, 771-774 (1998).
- Lendon, C., Ashall, F. & Goate, A. Exploring the etiology of Alzheimer's disease using molecular genetics. *J. Am. Med. Assoc.* **277**, 825-831 (1997).
- Levitán, D. & Greenwald, I. Effects of SEL-12 presenilin on LIN-12 localization and function in *Caenorhabditis elegans*. *Development* **125**, 3599-3606 (1998).
- De Strooper, B. et al. Production of intracellular amyloid-containing fragments in hippocampal neurons expressing human amyloid precursor protein and protection against amyloidogenesis by subtle amino acid substitutions in the rodent sequence. *EMBO J.* **14**, 4932-4938 (1995).
- Kopan, R., Schroeter, E. H., Weintraub, H. & Nye, J. S. Signal transduction by activated mNotch: importance of proteolytic processing and its regulation by the extracellular domain. *Proc. Natl Acad. Sci. USA* **93**, 1683-1688 (1996).
- Tischer, E. & Cordell, B. Beta-amyloid precursor protein. Location of transmembrane domain and specificity of gamma-secretase cleavage. *J. Biol. Chem.* **271**, 21914-21919 (1996).
- Brown, M. S. & Goldstein, J. L. The SREBP pathway: regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor. *Cell* **89**, 331-340 (1997).
- Ross, S. L. et al. Amyloid precursor protein processing in sterol regulatory element-binding protein site 2 protease-deficient Chinese hamster ovary cells. *J. Biol. Chem.* **273**, 15309-15312 (1998).
- Wolfe, M. S. et al. A substrate-based difluoro ketone selectively inhibits Alzheimer's gamma-secretase activity. *J. Med. Chem.* **41**, 6-9 (1998).

18. Higaki, J., Quon, D., Zhong, Z. & Cordell, B. Inhibition of beta-amyloid formation identifies proteolytic precursors and subcellular site of catabolism. *Neuron* **14**, 651–659 (1995).
19. Citron, M. *et al.* Evidence that the 42- and 40-amino acid forms of amyloid beta protein are generated from the beta-amyloid precursor protein by different protease activities. *Proc. Natl Acad. Sci. USA* **93**, 13170–13175 (1996).
20. Klafki, H., Abramowski, D., Swoboda, R., Paganetti, P. A. & Staufenbiel, M. The carboxyl termini of beta-amyloid peptides 1–40 and 1–42 are generated by distinct gamma-secretase activities. *J. Biol. Chem.* **271**, 28655–28659 (1996).
21. Yamazaki, T., Haass, C., Saido, T. C., Omura, S. & Ihara, Y. Specific increase in amyloid beta-protein 42 secretion ratio by calpain inhibition. *Biochemistry* **36**, 8377–8383 (1997).
22. Wolfe, M. S. *et al.* Peptidomimetic probes and molecular modelling suggest Alzheimer's γ -secretase is an intra-membrane cleaving aspartyl protease. *Biochemistry* (in the press).
23. Ray, W. J. *et al.* Evidence for a physical interaction between presenilin and Notch. *Proc. Natl Acad. Sci. USA* **96**, 3263–3268 (1999).
24. Weidemann, A. *et al.* Formation of stable complexes between two Alzheimer's disease gene products: presenilin-2 and beta-amyloid precursor protein. *Nature Med.* **3**, 328–332 (1997).
25. Xia, W., Zhang, J., Perez, R., Koo, E. H. & Selkoe, D. J. Interaction between amyloid precursor protein and presenilins in mammalian cells: implications for the pathogenesis of Alzheimer disease. *Proc. Natl Acad. Sci. USA* **94**, 8208–8213 (1997).
26. Saffig, P. & de Strooper, B. Downregulation of PS1 expression in neurons decreases beta-amyloid production: a biochemical link between the two major familial Alzheimer's disease genes. *Mol. Psychiat.* **3**, 287–289 (1998).
27. Rawson, R. B. *et al.* Complementation cloning of S2P, a gene encoding a putative metalloprotease required for intramembrane cleavage of SREBPs. *Mol. Cell* **1**, 47–57 (1997).
28. Varnum-Finney, B. *et al.* The Notch ligand, Jagged-1, influences the development of primitive hematopoietic precursor cells. *Blood* **91**, 4084–4091 (1998).
29. Robey, E. & Fowlkes, B. J. The $\alpha\beta$ versus $\alpha\delta$ T-cell lineage choice. *Curr. Opin. Immunol.* **10**, 181–187 (1998).
30. Chan, Y. M. & Jan, Y. N. Roles for proteolysis and trafficking in notch maturation and signal transduction. *Cell* **94**, 423–426 (1998).

Acknowledgements. This work was supported by the FWO-Vlaanderen, the Human Frontier of Science Program, the Flemish Institute for Biotechnology (VIB) and the K. U. Leuven. B.D.S., W.A., P.C. and K.C. are researchers of the FWO. R.K., M.S.W. and A.G. are supported by the NIH; W.J.R. is supported by a fellowship from the KECK foundation; J.S.M. is supported in part by the Markey Special Emphasis Pathway in Human Pathobiology; P.S. is supported by the Deutsche Forschungsgemeinschaft. We thank A. Roebroek and J. Gordon for helpful comments.

Correspondence and requests for materials should be addressed to B.D.S. (e-mail: Bart.Destrooper@med.kuleuven.ac.be) or R.K. (e-mail: Kopan@Pharmsun.Wustl.Edu).

Presenilin is required for activity and nuclear access of Notch in *Drosophila*

Gary Struhl^{*†} & Iva Greenwald^{‡‡}

Departments of ^{*}Genetics and Development, and [‡]Biochemistry and Molecular Biophysics, and [†]Howard Hughes Medical Institute, Columbia University, College of Physicians and Surgeons, New York, New York 10032, USA

Presenilins are membrane proteins with multiple transmembrane domains that are thought to contribute to the development of Alzheimer's disease by affecting the processing of β -amyloid precursor protein¹. Presenilins also facilitate the activity of transmembrane receptors of the LIN-12/Notch family^{2–5}. After ligand-induced processing, the intracellular domain of LIN-12/Notch can enter the nucleus and participate in the transcriptional control of downstream target genes^{6–9}. Here we show that null mutations in the *Drosophila Presenilin* gene abolish Notch signal transduction and prevent its intracellular domain from entering the nucleus. Furthermore, we provide evidence that presenilin is required for the proteolytic release of the intracellular domain from the membrane following activation of Notch by ligand.

β -Amyloid precursor protein (β -APP) is a transmembrane protein that travels by way of the endoplasmic reticulum and Golgi to the cell surface and undergoes proteolytic processing (reviewed in ref. 1). A set of β -amyloid (A β) peptides are generated from β -APP by proteases known as the β - and γ -secretases. The β -secretase cleavage occurs in the extracellular domain and the heterogeneous γ -secretase cleavages in the transmembrane domain. Dominant mutations in either of two human *Presenilin* genes appear to cause Alzheimer's disease by increasing the amount of the A β 42(43) fragment that is produced. A null allele of mouse *Presenilin1* appears selectively to reduce γ -secretase activity¹⁰. These observations indicate that presenilin either stimulates the activity of γ -secretase, or is itself a component of γ -secretase¹⁰.

LIN-12/Notch proteins act as transmembrane cell-surface receptors for intercellular signals during development. It has been proposed that signal transduction involves cleavage and transport of the intracellular domain to the nucleus (reviewed in ref. 11), and results from experiments in *Drosophila*^{6,8,9} and mammalian cells⁷ strongly support this idea, indicating that cleavage occurs in or near the transmembrane domain. In mammalian cells, at least one




Figure 1 Presenilin activity is required for Notch activity. **a**, Hunchback (Hb) expression marks neuroblasts. Notch activity normally prevents more than one cell in a proneural cluster from segregating as a neuroblast. In *PS*⁻ embryos, clusters of neuroblasts form instead of single neuroblasts, generating longitudinal bands, rather than lines, of neuroblasts. Hb expression is shown just after the completion of germ-band extension [the dorsal (top) and ventral (bottom) aspects of each embryo are shown in different focal planes; anterior is to the left]. **b**, Single-minded (Sim) expression marks ventral midline cells¹⁹. Notch activity normally promotes differentiation of midline cells. In *PS*⁻ (and *N*⁻) embryos, relatively few of these cells arise, in contrast to wild-type embryos where they form a continuous two-cell wide column along the midline. Sim expression is shown dorsally, just after completion of germ-band extension. **c**, Right: a wing containing clones of *PS*⁻ cells, marked with the *mwh* mutation (the *mwh* marker is not visible at this magnification). Notch activity is normally required to transduce signals between the dorsal and ventral compartments at the wing margin which are necessary for the survival and growth of wing blade cells²⁰. *PS*⁻ clones that arise near the wing margin (asterisks) cause scalloping of the wing, as do *N*⁻ clones in this region²⁵. Clones of mutant cells at a distance from the boundary can survive and populate internal portions of the wing blade (one such clone is indicated by an open arrow and outlined with dashed lines). However, such clones are associated with vein thickening (filled arrow), also attributable to a failure in signal transduction by Notch²¹. *PS*⁻/*PS*⁻ embryos derived from *PS*⁻/*+* females can survive until the onset of pupation; however, the imaginal discs are abnormally small. **d**, Notch protein (N) accumulates to similar levels and is mainly associated with the plasma membrane in *PS*⁺ and *PS*⁻ embryos. Both embryos are derived from *PS*⁻ female germ cells and are shown ventro-laterally at the same stage of gastrulation with the presumptive Sim-expressing midline cells forming single cell columns flanking the ventral midline. Right: *PS*⁻/*PS*⁻ zygote; left: *PS*⁺/*+* zygote, marked by the striped expression of β -galactosidase from a paternally derived third chromosome carrying both a *PS*⁺ allele and a *ftz-lacZ* reporter transgene (seven stripes of β -galactosidase expression can be seen superimposed on the otherwise normal pattern of N-protein expression). Such paternally rescued embryos can develop into normal first-instar larvae and give rise to adults that appear phenotypically wild-type.